

Separability in Riemannian manifolds

BY SERGIO BENENTI

*Dipartimento di Matematica, Università di Torino, via Carlo Alberto 10, 10123
Torino, Italy*

A short account of the basic Riemannian structures underlying the separation of variables in the Hamilton-Jacobi equation of natural Hamiltonian systems.†

Keywords: Riemannian geometry, Hamilton-Jacobi equation, separation

1. Introduction

In the last years the study of the **separable systems** (whose Hamilton-Jacobi equations can be integrated by separation of variables) has known a remarkable development, also in relation with other kinds of integrable systems (bi-Hamiltonian systems, Lax systems). I think that an outline of the Riemannian background of this theory may be useful for specialists as well as for beginners. With the exception of the last section, we shall confine our discussion to the **orthogonal separable systems** (or **Stäckel systems**) and to a special class of Stäckel systems, here called **L-systems**. Some theorems presented here are new. For some of the recalled theorems a new shorter proof is given.

Let Q be a n -dimensional Riemannian manifold with generic local coordinates $\underline{q} = (q^i)$ and (contravariant) metric tensor $\mathbf{G} = (g^{ij})$, which we assume to be positive-definite, and let T^*Q be the cotangent bundle of Q , with canonical coordinates $(\underline{q}, \underline{p}) = (q^i, p_i)$. We shall deal with the additive separation of the Hamilton-Jacobi equations

$$G(\underline{q}, \underline{p}) = E, \quad H(\underline{q}, \underline{p}) = E, \quad p_i = \partial_i W,$$

where $G = \frac{1}{2} g^{ij} p_i p_j$ is the **geodesic Hamiltonian** on T^*Q , and $H = G + V = \frac{1}{2} g^{ij} p_i p_j + V$ is a **natural Hamiltonian**, being $V(\underline{q})$ the **potential energy**, a smooth function on Q canonically lifted to a function on T^*Q . A coordinate system \underline{q} is called **separable** if the geodesic Hamilton-Jacobi equation $G = E$ admits a **complete solution** of the form

$$W(\underline{q}, \underline{c}) = \sum_{i=1}^n W_i(q^i, \underline{c}), \quad \underline{c} = (c^i), \quad \det \left[\frac{\partial^2 W}{\partial q^i \partial c^j} \right] \neq 0. \quad (1.1)$$

† This paper has been submitted to Royal Society and accepted for publication in a special volume dedicated to the state of the art of the separation of variables. Unfortunately, this volume was never published due to death of the Editor, V. Kutznetsov. Download: <http://www2.dm.unito.it/~benenti/CP/SOV.htm>.

Such a solution is called **separated solution**. Note that in these definitions the presence of a set of n constants \underline{c} satisfying the **completeness condition** (1.1)₂ is fundamental. Note that here we consider only **natural canonical coordinates**, where \underline{q} are coordinates on the configuration manifold. Levi-Civita (1904) proved that a *Hamilton-Jacobi equation* $H(\underline{q}, \underline{p}) = E$ admits a separated solution (1.1) if and only if the differential equations (notation: $\partial_i = \partial/\partial q^i$, $\partial^i = \partial/\partial p_i$)

$$L_{ij}(H) \doteq \partial_i H \partial_j H \partial^i \partial^j H + \partial^i H \partial^j \partial_i \partial_j H - \partial_i H \partial^j H \partial^i \partial_j H - \partial^i H \partial_j H \partial_i \partial^j H = 0,$$

are identically satisfied. They are called the **separability conditions** or **separability equations of Levi-Civita**. They provide not only a simple method for testing if a coordinate system is separable or not, but also the basis for the geometrical (i.e., intrinsic) characterization of the separation. A first (and well known) example of application is the following: the Levi-Civita equations for a natural Hamiltonian, $L_{ij}(G + V) = 0$, are polynomial equations of fourth degree in the momenta \underline{p} , which must be identically satisfied for all admissible values of these variables. It is easy to note that the fourth-degree homogeneous part of these equations is $L_{ij}(G) = 0$. This means that: (i) the separation of the geodesic equation is a necessary condition for the separation of equation $G + V = E$; (ii) the study of the geodesic separation plays a prominent role, (iii) the above-given definition of *separable coordinates* makes sense.

A special but fundamental case in this theory is the **orthogonal separation**, where the coordinates are assumed to be orthogonal, $g^{ij} = 0$ for $i \neq j$. In this case, examined firstly by Stäckel (1893), later on by Levi-Civita (1904), Eisenhart (1934, 1949), and more recently by many authors, the Levi-Civita equations $L_{ij}(G) = 0$ are equivalent to equations

$$S_{ij}(g^{kk}) = 0, \quad i \neq j, \quad (1.2)$$

where $S_{ij}(\cdot)$ denote the **Stäckel operators** associated with an orthogonal metric (g^{ii}). For any smooth function V on Q , it is defined by

$$S_{ij}(V) \doteq \partial_i \partial_j V - \partial_i \ln g^{jj} \partial_j V - \partial_j \ln g^{ii} \partial_i V, \quad i \neq j.$$

The Levi-Civita equations $L_{ij}(G + V) = 0$ are equivalent to

$$S_{ij}(g^{kk}) = 0, \quad S_{ij}(V) = 0. \quad (1.3)$$

2. Killing tensors

As shown by Eisenhart (1934, 1939) (for the orthogonal case) and by Kalnins & Miller (1980, 1981), the geodesic separation is related to the existence of Killing vectors and Killing tensors of order two. In this section we recall the basic properties of these objects. The contravariant symmetric tensors $\mathbf{K} = (K^{i\dots j})$ on Q are in one-to-one correspondence with homogenous polynomials on T^*Q ,

$$\mathbf{K} = (K^{i\dots j}) \longleftrightarrow P_{\mathbf{K}} = P(\mathbf{K}) = K^{i\dots j} p_i \dots p_j.$$

For a tensor of order zero i.e., a function f on Q , we define $P_f \doteq f$, where f is canonically lifted to T^*Q (by constant values on the fibers). The space of these polynomial functions is closed w.r.to the canonical Poisson bracket

$$\{A, B\} \doteq \partial^i A \partial_i B - \partial^i B \partial_i A.$$

Hence, on the space of the symmetric contravariant tensors we define a Lie-algebra structure $[\cdot, \cdot]$ by setting

$$P([\mathbf{K}_1, \mathbf{K}_2]) = \{P(\mathbf{K}_1), P(\mathbf{K}_2)\},$$

and the **symmetric product** \odot by setting

$$P(\mathbf{K}_1 \odot \mathbf{K}_2) = P(\mathbf{K}_1) \cdot P(\mathbf{K}_2).$$

Note that all the above-given definitions do not depend on a metric tensor. If a metric tensor \mathbf{G} is present, then we say that \mathbf{K} is a **Killing tensor** (KT) if $P(\mathbf{K})$ is in involution with $P(\mathbf{G}) = 2G$,

$$\{P(\mathbf{K}), P(\mathbf{G})\} = 0 \iff [\mathbf{K}, \mathbf{G}] = 0. \quad (2.1)$$

This means that $P(\mathbf{K})$ is a first integral of the geodesic flow. In the special case of a function f , this definition is equivalent to $\nabla f = 0$ (by ∇f we denote the **gradient** of a function f). A vector field \mathbf{X} is a **Killing vector**, $[\mathbf{X}, \mathbf{G}] = 0$, if and only if its flow preserves the metric.

Let us consider the case of a symmetric 2-tensor \mathbf{K} . Since a metric tensor is present, the boldface object \mathbf{K} can be represented in components as a tensor of type (2, 0), (1, 1) and (0, 2), respectively, $\mathbf{K} = (K^{ij}) = (K_j^i) = (K_{ij})$.

As a symmetric tensor of type (1, 1), \mathbf{K} defines an endomorphism on the space $\mathcal{X}(Q)$ of the (smooth) vector fields on Q and an endomorphism on the space $\Phi^1(Q)$ of the (smooth) 1-forms on Q . We shall denote by \mathbf{KX} the vector field image of $\mathbf{X} \in \mathcal{X}(Q)$ by \mathbf{K} , and by $\mathbf{K}\phi$ the 1-form image of $\phi \in \Phi^1(Q)$ by \mathbf{K} . This means that $\mathbf{KX} = K_j^i X^j \partial_i$, $\mathbf{K}\phi = K_j^i \phi_i dq^j$. Note that the metric tensor \mathbf{G} coincide with the identity operator \mathbf{I} , whose (1, 1) components are given by the Kronecker symbol δ_j^i . Then a 2-tensor \mathbf{K} gives rise to eigenvalues, eigenvectors or eigenforms, according to equations $\mathbf{KX} = \rho\mathbf{X}$, $\mathbf{K}\phi = \rho\phi$. We recall that, in a positive-definite metric, (i) all symmetric tensors have real eigenvalues; (ii) the *algebraic multiplicity* of an eigenvalue ρ (i.e., its order as a root of the characteristic equation $\det(\mathbf{K} - \rho\mathbf{G}) = 0$) is equal to its *geometrical multiplicity* (i.e., the dimension of the space of the corresponding eigenvectors, or eigenforms); the eigenspaces corresponding to distinct eigenvalues are orthogonal. We shall denote by $\mathbf{K}_1\mathbf{K}_2$ the product of the two endomorphisms \mathbf{K}_1 and \mathbf{K}_2 ; in components $(\mathbf{K}_1\mathbf{K}_2)^{ij} = K_1^{ih}K_{2h}^j$. The **algebraic commutator** of two tensors will be denoted by

$$[[\mathbf{K}_1, \mathbf{K}_2]] \doteq \mathbf{K}_1\mathbf{K}_2 - \mathbf{K}_2\mathbf{K}_1.$$

If a symmetric 2-tensor \mathbf{K} can be diagonalized in orthogonal coordinates, $K^{ij} = 0$ for $i \neq j$, then $K^{ii} = \rho^i g^{ii}$, where (ρ^i) are the eigenvalues of \mathbf{K} . By writing the **Killing equation** (2.1) in these coordinates, we see that \mathbf{K} is a KT if and only if equations

$$\partial_i \rho^j = (\rho^i - \rho^j) \partial_i \ln g^{jj}. \quad (2.2)$$

are satisfied by the eigenvalues. These equations have been called **Killing-Eisenhart equations** in Benenti *et al.* (2002a), since they have been extensively used by Eisenhart (1949). However, they already appear in a paper of Levi-Civita (1896), p.285.

3. Killing-Stäckel spaces

Equations (2.2) can be interpreted as a linear system of n first-order partial differential equations in normal form, in the n unknown functions $\rho^i(q)$. It is remarkable the fact that the integrability conditions assume of the form

$$(\rho^i - \rho^j) S_{ij}(g^{kk}) = 0.$$

Then their link with the orthogonal separation is at once clear. A second, and even more remarkable property, is that the unknown functions ρ^i appear in the integrability conditions through their differences $\rho^i - \rho^j$. This means that *if the system (2.2) admits a solution such that $\rho^i \neq \rho^j$, then it is completely integrable*. Note that for a linear system the converse is always true. Going back to the orthogonal separability conditions (1.2) we can immediately conclude that: (I) *a system of orthogonal coordinates is separable if and only if there exists a KT which is diagonalized in these coordinates and which has pointwise simple eigenvalues*. Furthermore, since system (2.2) is linear, if it is completely integrable then it admits a n -dimensional space of solutions (and conversely). As a consequence: (II) *a Killing tensor \mathbf{K} which has simple eigenvalues and is diagonalized in orthogonal coordinates generates a n -dimensional space \mathcal{K} of Killing tensors which are all diagonalized in the same coordinates*. Such a space will be called **Killing-Stäckel space** (KS-space). In the space of direct sums of Killing tensors, $\mathbf{K} = c \oplus \mathbf{K}_1 \oplus \mathbf{K}_2 \oplus \dots \oplus \mathbf{K}_n \oplus \dots$ endowed with the Lie bracket $[\cdot, \cdot]$ defined above, a KS-space, which is made of elements $0 \oplus 0 \oplus \mathbf{K}_2 \oplus 0 \oplus \dots$, is an involutive subalgebra. For this reason it has also been called *Killing-Stäckel algebra* in Benenti *et al.* (2002a).

Three remarks are in order: (i) *the metric tensor belongs to any KS-space* (indeed, $\rho^i = 1$ is a trivial solution of system (2.2)); (ii) *if two KS-spaces have an element with simple eigenvalues in common, then they coincide*; (iii) *all elements of a KS-space are in involution* (if equations (2.2) are satisfied for two tensors \mathbf{K}_1 and \mathbf{K}_2 , then $\{P(\mathbf{K}_1), P(\mathbf{K}_2)\} = 0$).

All the above properties have a local character and are related to a coordinate system. We remark, however, that they are more precisely related to an equivalence class of orthogonal systems, being equivalent two systems of coordinates q and q' simply related by a **separated transformation** or a **rescaling**: $q^i = q^i(q'^i)$.

We look for a coordinate-free description of all this matter. To this end we recall some basic concepts.

A **frame** on a differentiable manifold Q (not necessarily Riemannian) is a set of vector fields (\mathbf{X}_i) which form a basis of the tangent space $T_q Q$ at each point q of their domain of definition. In general, frames exist only locally. Global frames are defined if and only if the manifold is parallelizable i.e., when $TQ \simeq Q \times \mathbb{R}^n$. Two frames (\mathbf{X}_i) and (\mathbf{X}'_i) are said to be **equivalent** if there are nowhere vanishing functions f_i such that $\mathbf{X}_i = f_i \mathbf{X}'_i$. A frame is called **holonomic** or **integrable** if it is equivalent to a **natural frame** (∂_i) associated with coordinates (q^i) . A basic property is (cf. Schouten 1954 and Benenti *et al.* 2002a)

Theorem 3.1. *The three following conditions are equivalent: (i) the frame (\mathbf{X}_i) is holonomic, (ii) for each pair of indices (i, j) the distribution spanned by the vectors \mathbf{X}_i and \mathbf{X}_j is completely integrable, (iii) for each index i the distribution spanned by the $n - 1$ vectors \mathbf{X}_j for $j \neq i$ is completely integrable.*

On a Riemannian manifold a vector field \mathbf{X} is called **normal** if it is *orthogonally integrable* or *surface forming* i.e., if it is orthogonal to a family of hypersurfaces. In a positive-definite metric a symmetric tensor \mathbf{K} with simple eigenvalues and normal eigenvectors gives rise to an equivalence class of holonomic orthogonal frames hence, to an equivalence class of orthogonal coordinates. Then we get the following simple intrinsic characterization of the orthogonal geodesic separation (Kalnins & Miller 1980, corollary 4, §3; see also Benenti 1993):

Theorem 3.2. *The geodesic Hamilton-Jacobi equation is separable in orthogonal coordinates if and only if there exists a Killing 2-tensor with simple eigenvalues and normal eigenvectors.*

A KT having these properties will be called a **characteristic Killing tensor** (ChKT). The eigenvectors generate a family of n orthogonal foliations of manifolds of codimension 1, which we call **Stäckel web**. Any coordinate system (q^i) such that the web is locally described by equations $q^i = \text{const.}$ (this is equivalent to say that dq^i are eigenforms of the ChKT), is separable.

In accordance with the remarks above, a ChKT generates a KS-space. A suitable coordinate-independent definition of this concept is the following: a **Stäckel space** on a Riemannian manifold Q_n is a n -dimensional linear space \mathcal{K}_n of Killing 2-tensors whose elements (a) *commute as linear operators*, $[[\mathbf{K}_1, \mathbf{K}_2]] = 0$, and (b) *are in involution*, $[\mathbf{K}_1, \mathbf{K}_2] = 0$. Indeed, in the algebraic realm it can be proved that in such a space there exists an element with pointwise distinct eigenvalues (in the neighborhood of any given point of the domain of definition of \mathcal{K}_n); as a consequence, the commutation relation (a), applied to such a tensor \mathbf{K}_1 , shows that all elements have common eigenvectors. Furthermore, from (b) it follows that

Theorem 3.3 (Kalnins & Miller). *If n independent KT's in involution have the same eigenvectors, then these eigenvectors are normal.*

This very remarkable property was firstly discovered by Kalnins & Miller (1980) (proof of theorem 6, §3). However, it is also remarkable the fact that in this last theorem the assumption that the independent tensors are KT's is redundant. In fact, it can be proved that

Theorem 3.4. *An orthogonal frame made of common eigenvectors of n independent symmetric 2-tensors in involution is holonomic (the eigenvectors are normal).*

For a detailed discussion and proof see Benenti *et al.* (2002a). Hence, we have a second intrinsic characterization of the orthogonal geodesic separation (Kalnins & Miller 1980, theorem 6, §3; Shapovalov 1981; note that in theorem 6 of Kalnins & Miller 1980 item (4) is redundant):

Theorem 3.5. *The geodesic Hamilton-Jacobi equation is separable in orthogonal coordinates if and only if the Riemannian manifold admits a KS-space i.e., a n -dimensional linear space \mathcal{K} of Killing tensors commuting as linear operators and in involution.*

In the applications, one of these last two theorems can be used according to the convenience. In applying theorem 3.2 we have the advantage of dealing with a single KT, but difficulties may arise in testing if it has simple eigenvalues and

normal eigenvectors. However, for solving this problem we can use the following two effective criteria:

Theorem 3.6. *A $(1, 1)$ tensor \mathbf{K} has distinct eigenvalues if and only if*

$$D \doteq \begin{vmatrix} n & S_1 & \dots & S_{n-1} \\ S_1 & S_2 & \dots & S_n \\ \vdots & \vdots & \dots & \vdots \\ S_{n-1} & S_n & \dots & S_{2n-2} \end{vmatrix} \neq 0, \quad S_p \doteq \text{tr}(\mathbf{K}^p).$$

Here \mathbf{K}^p is the power $p = 0, 1, 2, \dots$ of the linear mapping \mathbf{K} . This theorem is a consequence of a classical theorem of Sylvester about the **discriminant** D of an algebraic equation, here applied to the characteristic equation of \mathbf{K} .

Theorem 3.7 (Haantjes, 1955). *A symmetric tensor \mathbf{K} with simple eigenvalues has normal eigenvectors if and only if*

$$H_{ab}^h K_i^a K_j^b + 2 H_{a[i}^b K_{j]}^a K_b^h + H_{ij}^a K_b^h K_a^b = 0,$$

where H is the Nijenhuis torsion of \mathbf{K} ,

$$H_{ij}^h(\mathbf{K}) \doteq 2 K_{[i}^a \partial_{|a|} K_{j]}^h - 2 K_a^h \partial_{[i} K_{j]}^a.$$

This is a special case of a more general theorem due to Haantjes (1955) (see also Schouten (1954), p. 248).

As it will be seen in §7, it is interesting the case of a **torsionless** tensor: $\mathbf{H}(\mathbf{K}) = 0$. We shall apply the following

Theorem 3.8 (Nijenhuis, 1951). *A symmetric tensor \mathbf{K} with simple eigenvalues ρ^i is torsionless if and only if it has normal eigenvectors \mathbf{X}_i such that*

$$\mathbf{X}_i \rho^j = 0, \quad i \neq j.$$

This means that each eigenvalue ρ^j is constant on the hypersurfaces orthogonal to the corresponding eigenvector \mathbf{X}_j . It is worthwhile to observe that

Theorem 3.9. *A torsionless KT with simple eigenvalues is necessarily a constant KT on a flat Riemannian manifold.*

Proof. If $\mathbf{H}(\mathbf{K}) = 0$, then equations (2.2) implies

$$0 = \partial_i \rho^j = (\rho^i - \rho^j) \partial_i \ln g^{jj}, \quad i \neq j, \quad \partial_i \rho^i = 0.$$

This shows that $\rho^i = \text{const.}$ and that $\partial_i g^{jj} = 0$ for $i \neq j$. This last condition means that g^{jj} is a function of q^j only. In this case, up to a change of scale of the coordinates, we can consider $g^{jj} = \text{const.}$ \square

This is a case considered in Bruce, McLenaghan & Smirnov 2001.

Going back to theorem 3.5, it is interesting to make a comparison with the intrinsic characterizations of the geodesic orthogonal separation due to Eisenhart (1934, 1949) and Woodhouse (1975). In the **Eisenhart theorem** (Eisenhart, 1934; p. 289) the necessary and sufficient conditions for the orthogonal separation are:

(i) the existence of $n - 1$ independent Killing tensors $\mathbf{K}_1, \dots, \mathbf{K}_{n-1}$ with normal common eigenvectors and such that (ii) for *each* of these tensors the eigenvalues are simple and (iii) for any pair (i, α) of fixed indices ($\alpha = 2, \dots, n$, $i = 1, \dots, n$) the square matrices $\|\rho_i^\alpha - \rho_j^\alpha\|$ (with $j \neq i$) are regular. In the Eisenhart notation, ρ_i^α are the eigenvalues of \mathbf{K}_α . Condition (i) should be replaced by the existence of $n - 1$ Killing tensors such that $\mathbf{G}, \mathbf{K}_1, \dots, \mathbf{K}_{n-1}$ are independent. Then theorem 3.5 shows that conditions (ii) and (iii) are redundant. In Woodhouse 1975 (theorem 4.2) the $n - 1$ KT's are assumed to be in involution and with common *closed* eigenforms. Theorem 3.3 shows that the requirement *closed* is redundant, since it is equivalent to the normality of the eigenvectors.

4. The orthogonal separation of a natural Hamiltonian

With each symmetric 2-tensor \mathbf{K} and a function U on Q we associate the function $F = \frac{1}{2} P_{\mathbf{K}} + U$ on T^*Q . We observe that F is a *first integral* of $H = G + V$, $\{H, F\} = 0$, if and only if

$$\{G, P_{\mathbf{K}}\} = 0, \quad dU = \mathbf{K} dV.$$

The first equation means that \mathbf{K} is a Killing tensor. If it is a ChKT, then, in any orthogonal coordinate system determined by its eigenvectors, the second equation is equivalent to $\partial_i U = \rho^i \partial_i V$. Due to the fundamental equations (2.2), the integrability conditions of these equations assume the form $\partial_j \partial_i U - \partial_i \partial_j U = (\rho^i - \rho^j) S_{ij}(V) = 0$. This proves

Theorem 4.1. *If \mathbf{K} is a symmetric 2-tensor with simple eigenvalues and normal eigenvectors, then F is a first integral of $G + V$ if and only if \mathbf{K} is a Killing tensor and $S_{ij}(V) = 0$ in any orthogonal system of coordinates generated by the eigenvectors.*

Since the existence of coordinates is a local matter, condition $dU = \mathbf{K} dV$ can be replaced by $d(\mathbf{K} dV) = 0$. Thus, by recalling theorem 3.2 and the remarks at the end of §1, we find (Benenti, 1993)

Theorem 4.2. *The Hamilton-Jacobi equation $G + V = E$ is separable in orthogonal coordinates if and only if there exists a Killing 2-tensor \mathbf{K} with simple eigenvalues and normal eigenvectors such that*

$$d(\mathbf{K} dV) = 0. \tag{4.1}$$

This equation has been called **characteristic equation** of a **separable potential** V . We observe that for $n = 2$ any vector field is normal. Since it can be proved that on a two-dimensional manifold the separation always occurs in orthogonal coordinates (Levi-Civita 1904), we get

Theorem 4.3. *On a two-dimensional Riemannian manifold the Hamilton-Jacobi equation $G + V = E$ is separable if and only if there exists a (non-trivial) quadratic first integral.*

This is the extension to a two-dimensional manifold of the so-called Bertrand-Darboux-Whittaker theorem for the Euclidean plane \mathbb{E}_2 (Whittaker 1937, §§152, 153; Ankiewicz & Pask 1983).

When written in Cartesian coordinates (on a flat manifold or on a Euclidean n -space, equation (4.1) gives rise to the so-called **Bertrand-Darboux** (BD) equations. If we know the form of all characteristic tensors of a manifold, then equation (4.1), written in any coordinate system (even not separable), provides an effective criterion for the separability of the Hamilton-Jacobi equation. This criterion have been applied for instance in the study of the super-separability of the inverse-square three-dimensional Calogero system (Benenti *et al.*, 2000). In Benenti *et al.* (2000) we can find the intrinsic (i.e., "boldface") expressions of all the characteristic Killing 2-tensors in the Euclidean three-space \mathbb{E}_3 , so that this criterion is ready to be used for any potential V . For a general \mathbb{E}_n , the basic ChKT's and the corresponding BD equations have been determined in Marshall & Wojciechowski 1988 and in Benenti 1992*b*, 1993. This analysis has been completed by Waksjö 2000. In his thesis he presents an effective general criterion for the separability of a potential V in the Euclidean n -space. See also Waksjö & Rauch-Wojciechowski 2003. Other separability criteria can be based on the analysis of the fundamental invariants of spaces of Killing tensors under the action of isometry groups and the method of moving frames (see, also for related references, McLenaghan & Smirnov 2002, Deeley *et al.* 2004).

If in \mathbb{E}_n we take

$$\mathbf{K} = \text{tr}(\mathbf{L}) \mathbf{G} - \mathbf{L}, \quad \mathbf{L} \doteq \mathbf{A} + m \mathbf{r} \otimes \mathbf{r} + \mathbf{w} \odot \mathbf{r}, \quad (4.2)$$

where \mathbf{A} is symmetric and constant, $m \in \mathbb{R}$, \mathbf{w} is a constant vector, and \mathbf{r} is the vector representing the generic point, then in Cartesian coordinates (x^i) equation (4.1) yields the Bertrand-Darboux equations (2.2) of Rauch-Wojciechowski 2004, in this volume, or equations (3.25) of Waksjö 2000. The correspondence of notation is the following: $\mathbf{A} = (\gamma_{ij})$, $m = \alpha$, $\mathbf{w} = (2\beta_i)$.

For $\mathbf{w} = 0$ we get the BD equations for the separation in elliptic coordinates centered at the origin. For $m = 0$ we get the BD equations for the separation in parabolic coordinates centered at the point P where $\mathbf{L}_P(\mathbf{w})$ (the existence of such a point is proved in Benenti 1992*b*). For $\mathbf{w} = 0$ and $m = 0$ we have the separation in Cartesian coordinates. In the remaining case $\mathbf{w} \neq 0$ and $m \neq 0$, we have the separation in elliptic coordinates centered in the point $\mathbf{c} = -\frac{1}{2m} \mathbf{w}$ (see below).

It must be emphasized that in the characteristic equation (4.1) for a separable potential V the eigenvalues of \mathbf{K} must be simple (outside a **singular set**). For the present case we have

Theorem 4.4. *The tensor \mathbf{K} defined by (4.2) has simple eigenvalues if and only if for $m = 0$ the eigenvalues of \mathbf{A} are simple and for $m \neq 0$ the eigenvalues of $\mathbf{A} - \frac{1}{4m} \mathbf{w} \otimes \mathbf{w}$ are simple.*

This is (a slightly modified version of) a theorem of Waksjö 2000, p. 45. We give here an alternative proof.

Proof. The tensor \mathbf{K} has simple eigenvalues if and only if \mathbf{L} has simple eigenvalues. For $m = 0$ we have $\mathbf{L} = \mathbf{A} + \mathbf{w} \odot \mathbf{r}$; we are in the case of the parabolic web (Benenti 1992*b*) and \mathbf{L} has simple eigenvalues if and only if \mathbf{A} has simple (constant) eigenvalues. For $m \neq 0$, let us change the origin by considering the transformation $\mathbf{r} = \mathbf{c} + \mathbf{r}'$, where \mathbf{c} is a constant vector. We obtain

$$\mathbf{L} = \mathbf{A} + m (\mathbf{r}' \otimes \mathbf{r}' + 2 \mathbf{r}' \odot \mathbf{c} + \mathbf{c} \otimes \mathbf{c}) + \mathbf{w} \odot \mathbf{c} + \mathbf{w} \odot \mathbf{r}'.$$

If we take $\mathbf{c} = -\frac{1}{2m} \mathbf{w}$, then

$$\mathbf{L} = \mathbf{A} - \frac{1}{4m} \mathbf{w} \otimes \mathbf{w} + m \mathbf{r}' \otimes \mathbf{r}' = \mathbf{A}' + m \mathbf{r}' \otimes \mathbf{r}'.$$

We know that for a symmetric tensor of the kind $\mathbf{A}' + \mathbf{r}' \otimes \mathbf{r}'$ the eigenvalues (u^i) are the roots of equation

$$\sum_i \frac{x_i^2}{u - a^i} = \frac{1}{m},$$

where (x_i) are Cartesian coordinates and (a^i) are the constant eigenvalues of \mathbf{A}' . This equation is equivalent to

$$m \sum_i x_i^2 \prod_{k \neq i} (u - a^k) - \prod_j (u - a^j) = 0. \quad (4.3)$$

If (a^i) are simple, then also (u^i) are simple, since $a^1 < u^1 < a^2 < u^2 < \dots < u^{n-1} < a^n < u^n$. If (a^i) are not all simple, for instance $a_1 = a_2$, then equation (4.3) has a double root $u = a^1$. \square

The tensors \mathbf{K} and \mathbf{L} have a mechanical meaning: they were introduced in Benenti (1992b) as the **inertia tensor** and **planar inertia tensor** of a set of massive points (including, this is important, negative masses) in \mathbb{E}_n . The parameter m is just the total mass (it may be 0). Indeed, it is a remarkable fact that an inertia tensor is a KT. This interpretation is of help in the problem of finding the intrinsic expressions of all the ChKT's of \mathbb{E}_n (see also Marshall & Wojciechowski 1988).

Notation. A matrix of the kind $\mathbf{A} + \mathbf{r} \otimes \mathbf{r}$ has been used by Moser (1981) for constructing a Lax pair for the geodesic flow of an asymmetric ellipsoid. For this reason it was denoted by \mathbf{L} in Benenti (1992b) (indeed, in analogy with the Lax method, starting from \mathbf{L} we can construct by a pure algebraic process, a complete system of first integrals in involution). There were other two reasons which suggested this notation: (i) \mathbf{L} is a letter adjacent to \mathbf{K} , and this is appropriate because a tensor \mathbf{L} generates a tensor \mathbf{K} according to (4.2); (ii) \mathbf{L} stands for *Levi-Civita*, and indeed the orthogonal metric associated with \mathbf{L} was firstly introduced by Levi-Civita (1896) (see the end of §7).

5. First integrals associated with the orthogonal separation

Going back to the characteristic equation (4.1) we recall that at the beginning of §3 we observed that a characteristic Killing tensor (like that appearing in equation (4.1)) generates a KS-space \mathcal{K} . It is remarkable the fact that

Theorem 5.1. *If $(\mathbf{K}_a) = (\mathbf{K}_0, \mathbf{K}_2, \dots, \mathbf{K}_{n-1})$ is a basis of \mathcal{K} (\mathbf{G} and \mathbf{K} may belong to this basis) then locally there exist functions V_a such that*

$$H_a \doteq \frac{1}{2} P_{\mathbf{K}_a} + V_a \quad (5.1)$$

are independent first integrals in involution.

An easy proof can be found in Benenti (1993). Indeed, it can be shown that equation (4.1) implies $d(\mathbf{K}_a dV) = 0$ for each index a . This implies that a ChKT

\mathbf{K} and a separable potential V generates a n -dimensional space $\mathcal{H}(\mathbf{K}, V)$ of first integrals in involution,

$$H_{\mathbf{K}} = \frac{1}{2} P_{\mathbf{K}} + V_{\mathbf{K}}, \quad \mathbf{K} \in \mathcal{K}, \quad (5.2)$$

and that the associated potentials $V_{\mathbf{K}}$ can be determined by integrating the closed 1-forms $\mathbf{K}_a dV$. We observe that there are separable systems (an example is the three-body Calogero system, Benenti *et al.* 2000) in which this integration can be avoided and replaced by an algebraic process.

We remark that, if we know a basis of a KS-space, then for testing if a potential V is separable it is sufficient to verify that equation (4.1) is satisfied for an element of this space with simple eigenvalues. When the answer is affirmative, then a complete set of integrals in involution can be determined by integrating the closed 1-forms $\mathbf{K}_a dV$.

It is well known that the orthogonal (as well as the non-orthogonal separation) is related to Stäckel matrices. A **Stäckel matrix** in the n variables (q^i) is a regular $n \times n$ matrix $\mathbf{S} = [\varphi_i^{(a)}]$ of functions $\varphi_i^{(a)}$ depending on the variable q^i corresponding to the lower index only. We denote by $[\varphi_{(a)}^i]$ the inverse matrix. The original **Stäckel theorem** asserts that *an orthogonal coordinate system (q^i) is separable if and only if there exists a Stäckel matrix such that $g^{ii} = \varphi_{(0)}^i$. In this case, (i) the diagonalized tensors $K_a^{ii} = \varphi_{(a)}^i$ are the basis of a Stäckel space, (ii) all separable potentials V have the form $V = \phi_i(q^i) \varphi_{(0)}^i$, where ϕ_i is a function of the corresponding coordinate q^i only; (iii) a basis H_a of the space of first integrals in involution is given by (5.1) with $V_a = \phi_i \varphi_{(a)}^i$.*

6. Conformal Killing tensors

As remarked above, the existence of a KT \mathbf{K} with simple eigenvalues and normal eigenvectors is a necessary and sufficient condition for the existence of a KS-space \mathcal{K} i.e., of a n -dimensional linear space of KT's with common normal eigenvectors (and, consequently, in involution). The following question arises: *is it possible to construct a basis of the space \mathcal{K} by a coordinate independent algebraic procedure, starting from \mathbf{K} ?* Note that this problem can be solved (in principle) by integrating the linear differential system (2.2), if we know a separable coordinate system.

As illustrated in Benenti (1992b), the answer is affirmative at least for special kinds of Stäckel systems. In the next sections we shall revisit this matter, by proposing new definitions and theorems. To this end, we need to recall some basic definitions and properties concerning conformal Killing tensors.

A **conformal Killing tensor** (CKT) on a Riemannian manifold Q_n is a symmetric tensor \mathbf{L} of order l satisfying equation $\{P_{\mathbf{L}}, P_{\mathbf{G}}\} = P_{\mathbf{X}} P_{\mathbf{G}}$, where \mathbf{X} is a suitable symmetric tensor of order $l - 1$. Since we are interested in CKT's of order two, we write this equation in the form

$$\{P_{\mathbf{L}}, P_{\mathbf{G}}\} = -2 P_{\mathbf{C}} P_{\mathbf{G}}, \quad (6.1)$$

where \mathbf{C} is a vector field which we call **associated with \mathbf{L}** (also denoted by $\mathbf{C}(\mathbf{L})$).

A CKT is said to be **of gradient-type** (GCKT) if $\mathbf{C} = \nabla f$. An example of GCKT is $f\mathbf{G}$. In this case we have $\mathbf{C} = \nabla f$. Indeed, $\{P_{f\mathbf{G}}, P_{\mathbf{G}}\} = \{f P_{\mathbf{G}}, P_{\mathbf{G}}\} = P_{\mathbf{G}}\{f, P_{\mathbf{G}}\} = -2g^{ij} p_j \partial_i f P_{\mathbf{G}}$. A KT is obviously a GCKT with $\mathbf{C} = 0$.

Proposition 6.1. *A CKT \mathbf{L} is of gradient-type with $\mathbf{C} = \nabla f$ if and only if $\mathbf{K} = f\mathbf{G} - \mathbf{L}$ is a Killing tensor.*

Proof. $\{P_{\mathbf{L}} - fP_{\mathbf{G}}, P_{\mathbf{G}}\} = \{P_{\mathbf{L}}, P_{\mathbf{G}}\} - P_{\mathbf{G}}\{f, P_{\mathbf{G}}\} = -2P_{\mathbf{C}}P_{\mathbf{G}} + 2P_{\mathbf{G}}P_{\nabla f}$. \square

A **conformal Killing tensor of trace-type \mathbf{L}** is a CKT for which $\mathbf{C} = \nabla \text{tr}(\mathbf{L})$. In this case, $\mathbf{K} = \text{tr}(\mathbf{L})\mathbf{G} - \mathbf{L}$ is a KT.

By considering in (6.1) $P_{\mathbf{G}} = g^{ii}p_i^2$ and $P_{\mathbf{L}} = u^i g^{ii}p_i^2$, we find

Theorem 6.2. *Assume that a symmetric 2-tensor \mathbf{L} is diagonalized in orthogonal coordinates, so that $L^{ii} = u^i g^{ii}$, $L^{ij} = 0$, $i \neq j$, where u^i are the eigenvalues. Then \mathbf{L} is a CKT if and only if the following equations are satisfied,*

$$\partial_i u^k = (u^i - u^k)\partial_i \ln g^{kk} + C_i, \quad C_i = \partial_i u^i. \quad (6.2)$$

7. L-tensors

Let us call **L-tensor** a (i) conformal Killing tensor \mathbf{L} with (ii) vanishing torsion and (iii) pointwise simple eigenvalues (u^i). The reasons for introducing such an object will be explained in the next section. In the present section we examine the basic properties of a L-tensor. Due to the vanishing of the torsion, there is an equivalence classes of orthogonal coordinates (q^i) in which this tensor is diagonalized and $\partial_i u^j = 0$ for $i \neq j$. We say that these coordinates are **associated** with \mathbf{L} . Since \mathbf{L} is a CKT, due to theorem 6.2 equations

$$\partial_i u^j = (u^i - u^j)\partial_i \ln g^{jj} + C_i = 0, \quad i \neq j, \quad C_i = \partial_i u^i \quad (7.1)$$

hold. For $\mathbf{C} = 0$ we find equations (2.2) of a KT.

Remark. In the above definition no assumption is made about the independence of the eigenvalues as functions on Q ; some of them may be constant (a criterion for the independence of the eigenvalues is given in theorem 9.2 below). This definition has to be compared with those given in Ibort *et al.* (2000), Błaszak (2003), Bolsinov & Matveev (2003), where \mathbf{L} is assumed to be a torsionless CKT of trace-type (and called *Benenti tensor*) and in Crampin (2003a, b), where \mathbf{L} is assumed to be a torsionless CKT with independent (i.e., coordinate-forming) eigenvalues (and called *special conformal Killing tensor*, see theorem 9.1 below). In all these papers the essential condition that \mathbf{L} has simple eigenvalues is missing (or understood). In fact it can be proved that

Theorem 7.1. *If an eigenvalue u^i of a torsionless CKT of trace-type is not simple, then it is constant.*

The proof of this theorem (here omitted) requires the use of the Haantjes theorem for a tensor with non-simple eigenvalues.

Theorem 7.2. *Let \mathbf{L} be a L-tensor with associated coordinates (q^i). Then: (i) Each eigenvalue u^i depends on the associated coordinate q^i only, $u^i = u^i(q^i)$. (ii) It is of trace-type, $\mathbf{C} = \nabla \text{tr}(\mathbf{L})$. (iii) In associated coordinates the metric has the form*

$$g^{kk} = \phi_k \prod_{i \neq k} \frac{1}{|u^i - u^k|}, \quad u^i = u^i(q^i), \quad \phi_k = \phi_k(q^k) > 0, \quad (7.2)$$

or, after a rescaling

$$g^{kk} = \prod_{i \neq k} \frac{1}{|u^i - u^k|}. \quad (7.3)$$

In both cases,

$$\partial_i \ln g^{kk} = \frac{\partial_i u^i}{u^k - u^i} \quad (i \neq k).$$

(We call **normal coordinates associated** with \mathbf{L} the orthogonal coordinates for which equations (7.3) hold). (iv) *The associated coordinates are separable.* (v) \mathbf{L} commutes with the Ricci tensor \mathbf{R} , $[\mathbf{L}, \mathbf{R}] = 0$ i.e., the Robertson condition is satisfied: in the associated coordinates, $R_{ij} = 0$ for $i \neq j$.

These properties are derived from Benenti 1992b, 1993. They follow from equations (7.1) and from the fundamental properties of the elementary symmetric polynomials (see the next section). From (7.2) it follows that the **contracted Christoffel symbols** $\Gamma_i = g^{hj} \Gamma_{hj,i}$ take the simple form $\Gamma_i = -\frac{1}{2} \phi'_k \phi_k$. Thus, in normal coordinates $\Gamma_i = 0$. Since in separable orthogonal coordinates $R_{ij} = \frac{3}{2} \partial_i \Gamma_j$, for $i \neq j$, the Robertson condition (v) is proved. Item (ii) (which follows from (7.1): $C_i = \partial_i u^i$ and $\partial_i u^j = 0$ for $i \neq j$ implies $C_i = \partial_i \sum_j u^j$) shows that in proposition 1, §1, of Crampin 2003a the assumption that (u^i) are functionally independent eigenvalues is not needed.

It is a remarkable fact that the metric (7.2) associated with \mathbf{L} is that of the **corresponding geodesics** found by Levi-Civita (1896): *there exists a metric $\bar{\mathbf{G}} = (\bar{g}^{ij})$ having the same (unparametrized) geodesics of a given metric $\mathbf{G} = (g^{ij})$ if and only if there are orthogonal coordinates in which the metric \mathbf{G} assumes the form (7.2).* It must be pointed out that this theorem holds under the assumption that the tensor $\bar{\mathbf{G}}$ has simple eigenvalues w.r.to \mathbf{G} . This matter has been recently analyzed by Bolsinov & Matveev (2003), and by Crampin (2003b). The metric (7.2) is a particular case of the orthogonal separable metric determined by Eisenhart (1949), App. 13, and characterized by the condition $R_{jikk} = 0$ for $i, j, k \neq$.

8. L-systems

Let $\sigma_a(\underline{u})$ denote the elementary symmetric polynomial of degree a of the n variables $\underline{u} = (u^i)$. Let σ_a^i and σ_a^{ij} be the functions obtained from σ_a by setting $u^i = 0$ and $u^j = 0$. Let us set

$$\sigma_0 = \sigma_0^i = \sigma_0^{ij} = 1, \quad \sigma_{-1} = \sigma_{-1}^i = \sigma_{-1}^{ij} = 0, \quad \sigma_n^{ij} = \sigma_n^{ij} = 0. \quad (8.1)$$

Then the following equations are satisfied (Benenti 1992b, §2):

$$\sigma_a = \sigma_a^i + u^i \sigma_{a-1}^i, \quad \sigma_a^j = \sigma_a^{ij} + u^i \sigma_{a-1}^{ij}, \quad \sigma_a^i - \sigma_a^j = (u^j - u^i) \sigma_{a-1}^{ij}. \quad (8.2)$$

$$\sum_i u^i \sigma_{a-1}^i = a \sigma_a, \quad \det[\sigma_a^i] = \prod_{j>i} (u^i - u^j). \quad (8.3)$$

$$\frac{\partial \sigma_a}{\partial u^i} = \sigma_{a-1}^i, \quad \frac{\partial \sigma_a^j}{\partial u^i} = \sigma_{a-1}^{ij}, \quad (u^j - u^i) \frac{\partial \sigma_a^j}{\partial u^i} = \sigma_a^i - \sigma_a^j. \quad (8.4)$$

If $u^i \neq u^j$ for $i \neq j$, then

$$\frac{\partial \sigma_a^j}{\partial u^i} = \frac{\sigma_a^i - \sigma_a^j}{u^j - u^i}, \quad \det[\sigma_a^i] \neq 0, \quad a = 0, 1, \dots, n-1, \quad i = 1, \dots, n. \quad (8.5)$$

It follows that for any coordinate system (q^i) ,

$$\partial_i \sigma_a^j = \partial_i u^h \frac{\partial \sigma_a^j}{\partial u^h} = \sum_{h \neq j} \partial_i u^h \sigma_{a-1}^{hj} = \sum_{h \neq j} \partial_i u^h \frac{\sigma_a^h - \sigma_a^j}{u^j - u^h}. \quad (8.6)$$

Theorem 8.1. *Let \mathbf{L} be a symmetric 2-tensor with eigenvalues (u^i) . The tensors $(\mathbf{K}_a) = (\mathbf{K}_0, \mathbf{K}_1, \dots, \mathbf{K}_{n-1})$ defined by*

$$\mathbf{K}_0 = \mathbf{G}, \quad \mathbf{K}_a = \frac{1}{a} \operatorname{tr}(\mathbf{K}_{a-1} \mathbf{L}) \mathbf{G} - \mathbf{K}_{a-1} \mathbf{L}, \quad a > 1, \quad (8.7)$$

or by

$$\mathbf{K}_a = \sum_{k=0}^a (-1)^k \sigma_{a-k} \mathbf{L}^k, \quad \mathbf{K}_a = \sigma_a \mathbf{G} - \mathbf{K}_{a-1} \mathbf{L} \quad (\mathbf{K}_{-1} = \mathbf{0}), \quad (8.8)$$

form a basis of a KS-space if and only if \mathbf{L} is a L-tensor.

Proof. (i) Assume that (\mathbf{K}_a) defined by (8.7) is a basis of a KS-space. Then they are linearly independent and there are orthogonal (separable) coordinates in which these tensors as well as \mathbf{L} are diagonalized and equations

$$\partial_i \rho_a^j = (\rho_a^i - \rho_a^j) \partial_i \ln g^{jj}, \quad \det [\rho_a^i] \neq 0, \quad (8.9)$$

hold, being ρ_a^i the eigenvalues of \mathbf{K}_a . Due to (8.7), these eigenvalues fulfill the recurrence relation $\rho_a^i = \frac{1}{a} \sum_k \rho_{a-1}^k u^k - \rho_{a-1}^i u^i$. On the other hand, from the first equations (8.3) and (8.2) we get $\frac{1}{a} \sum_k \sigma_{a-1}^k u^k - \sigma_{a-1}^i u^i = \sigma_a - \sigma_{a-1}^i u^i = \sigma_a^i$. This shows that

$$\rho_a^i = \sigma_a^i(u).$$

It follows that: (I) Due to the second equations (8.3) and (8.9), the eigenvalues u^i are simple. (II) Due to the first equation (8.2), the definition (8.7) implies the alternative definitions (8.8). (III) Due to the first equation (8.9), $\partial_i \rho_a^i = 0$ thus, $0 = \sum_{h \neq i} \partial_i u^h \sigma_{a-1}^{hi}$. Let us consider the case $i = 1$. We get the linear homogeneous system of $n - 1$ equations

$$\sum_{h>1} \partial_1 u^h \sigma_{a-1}^{h1} = 0, \quad a = 1, \dots, n - 1, \quad (8.10)$$

in the $n - 1$ unknown functions $\partial_1 u^h$, with $h = 2, \dots, n$. We can put $\sigma_{a-1}^{h1} = \tilde{\sigma}_b^h$, where $\tilde{\sigma}_b^h$, $b = 0, \dots, n - 2$, are the symmetric polynomials in the $n - 1$ variables (u^2, \dots, u^n) . In analogy with the third equation (8.4) we have $\det[\tilde{\sigma}_b^h] = \prod_{j>i>1} (u^i - u^j)$, thus $\det[\tilde{\sigma}_b^h] = \det[\sigma_{a-1}^{h1}] \neq 0$. It follows from (8.10) that $\partial_1 u^h = 0$ for all $h > 1$. In a similar way we prove that $\partial_i u^h = 0$ for all $h \neq i$. This shows that $\mathbf{H}(\mathbf{L}) = 0$. Finally, from $\mathbf{K}_1 = \operatorname{tr}(\mathbf{K}_1 \mathbf{L}) - \mathbf{L}$ we get

$$\mathbf{L} \doteq \frac{1}{n-1} \operatorname{tr}(\mathbf{K}_1) \mathbf{G} - \mathbf{K}_1. \quad (8.11)$$

Being \mathbf{K}_1 a KT, \mathbf{L} is a CKT. (ii) Conversely, assume that \mathbf{L} is a L-tensor. In coordinates associated with \mathbf{L} we have $\partial_i u^h = 0$ for all $h \neq i$, and moreover, $0 = (u^i - u^j) \partial_i \ln g^{jj} + \partial_i u^i$. By (8.6) we get

$$\partial_i \sigma_a^j = \partial_i u^i \frac{\sigma_a^i - \sigma_a^j}{u^j - u^i} = (\sigma_a^i - \sigma_a^j) \partial_i \ln g^{jj}.$$

Being $\rho_a^i = \sigma_a^i$, this shows that the tensors \mathbf{K}_a are Killing tensors. They are point-wise independent due to (8.3). \square

We call **L-system** any separable orthogonal system whose KS-space is generated by a L-tensor according to theorem 8.1. We call **L-web** any orthogonal web corresponding to a L-system.

Remarks.

(i) All the Killing tensors \mathbf{K}_a ($a \neq 0$) have simple eigenvalues. The sequence (8.7) was suggested by the analysis (Benenti 1992b) of the planar inertia tensor \mathbf{L} of an asymmetric massive body in the Euclidean n -space. The tensor \mathbf{K}_1 is the corresponding inertia tensor. Formula (8.8) shows that $\mathbf{K}_a = 0$ for $a > n - 1$, since for $a = n$ the right-hand side vanishes due to the Hamilton-Cayley theorem. (ii) Formula (8.7) is more effective than (8.8) since it does not require the knowledge of the eigenvalues of \mathbf{L} .

(ii) The sequence (8.7) is the same of that considered in Schouten (1954), p. 30, generated by a matrix \mathbf{L} (in Schouten denoted by \mathbf{P}). This sequence was firstly introduced by Fettis (1950) and Souriau (1950) for computing the eigenvectors of a matrix \mathbf{P} without solving systems of linear equations (if the eigenvalues are known and simple).

(iii) We observe (Schouten 1954) that the (1,1) tensor $\mathbf{Q}(x) = \text{cof}(\mathbf{L} - x\mathbf{G})$ is polynomial of degree $n - 1$ in x , whose coefficients, up to the sign, are the tensors \mathbf{K}_a defined in (8.8). We recall that the cofactor $\mathbf{A} = \text{cof}(\mathbf{A})$ of \mathbf{A} is defined by $\mathbf{A}\mathbf{A} = \mathbf{A}\mathbf{A} = \det(\mathbf{A})\mathbf{G}$. Hence, the Stäckel systems of the kind considered in the last two theorems are just the so-called **cofactor systems**, (cf. Błaszak & Ma 2003). In fact, they should be called **Levi-Civita systems** (so that "L-systems" is a good notation) since the separable metric (7.3) (but not the tensor \mathbf{L}) appears for the first time in Levi-Civita, 1896. In that paper it also shown that for such a metric the function

$$F(\underline{q}, \underline{\dot{q}}, c) \doteq \prod_{j \neq i} (w^j + c) |w^j - w^i| (\dot{q}^i)^2$$

is a first integral of the geodesics for all values of the parameter c . Since F is a polynomial in c of degree $n - 1$, its coefficients gives rise to n first integrals. These first integrals coincide, up to the sign and after the Legendre transformation, with the first integrals $P(\mathbf{K}_a)$.

(iv) Due to item (v) of theorem 7.2, a necessary condition for a Stäckel system to be a L-system is the Robertson condition: $\llbracket \mathbf{K}, \mathbf{R} \rrbracket = 0$ for a characteristic tensor \mathbf{K} (thus, for all elements of the KS-space). A criterion for testing if a Stäckel system is a L-system is the following.

Theorem 8.2. *A Stäckel system is a L-system if and only if in the corresponding KS-space there exists a characteristic tensor \mathbf{K}_1 such that the tensor \mathbf{L} defined by (8.11) is torsionless.*

Proof. (i) Assume that there exists a ChKT \mathbf{K}_1 such that the tensor \mathbf{L} defined by (8.11) is torsionless. Any tensor of the kind $f\mathbf{G} + \mathbf{K}$ is a CKT if \mathbf{K} is a KT. Since \mathbf{K}_1 has simple eigenvalues, also \mathbf{L} has simple eigenvalues. Then \mathbf{L} is a L-tensor. The tensors \mathbf{K}_a constructed by applying (8.7) form a KS-space \mathcal{K}_* which has \mathbf{K}_1 in common with the original KS-space \mathcal{K} . Thus, $\mathcal{K}_* = \mathcal{K}$ since two KS-spaces with a ChKT in common coincide. (ii) The converse is obvious. \square

About the *uniqueness* of a L-tensor generating a L-system, it can be proved that

Theorem 8.3. *Two L-tensors \mathbf{L} and $\tilde{\mathbf{L}}$ generates the same L-system if and only if $\tilde{\mathbf{L}} = a\mathbf{L} + b\mathbf{G}$, $a, b \in \mathbb{R}$, $a \neq 0$.*

In accordance with the remarks at the end of §5, a basis (\mathbf{K}_a) of a KS-space corresponds to the inverse of a Stäckel matrix $\mathbf{S}^{-1} = [\varphi_{(a)}^i]$, $\varphi_{(a)}^i = K_a^{ii}$. In the sequence (8.7) we have $\mathbf{K}_0 = \mathbf{G}$, then $\varphi_{(0)}^i = g^{ii}$ and $\varphi_{(a)}^i = K_a^{ii} = \sigma_a^i g^{ii}$. By using the formulas concerning the elementary symmetric function at the beginning of §8 it can be proved that

Theorem 8.4. *For a L-system the Stäckel matrix associated with the basis (8.7), $\mathbf{S} = [\varphi_{(a)}^i]$, assuming $u^1 < u^2 < \dots < u^n$, is of **alternating Vandermonde** type,*

$$\varphi_{(a)}^i = (-1)^{n-a+i} (u^i)^{n-a-1} \quad \begin{cases} a = 0, \dots, n-1 \\ i = 1, \dots, n \quad \text{index of row.} \end{cases} \quad (8.12)$$

For $n = 3$ we have (a index of row)

$$[\varphi_{(a)}^i] = \begin{bmatrix} g^{11} & g^{22} & g^{33} \\ (u^2 + u^3)g^{11} & (u^3 + u^1)g^{22} & (u^1 + u^2)g^{33} \\ u^2u^3g^{11} & u^3u^1g^{22} & u^1u^2g^{33} \end{bmatrix}$$

with $g^{11} = \frac{1}{(u^1-u^2)(u^1-u^3)}$, $g^{22} = \frac{1}{(u^2-u^3)(u^1-u^2)}$, $g^{33} = \frac{1}{(u^3-u^2)(u^3-u^1)}$. In accordance with theorem 8.11, the inverse matrix is

$$\mathbf{S} = [\varphi_{(a)}^i] = \begin{bmatrix} (u^1)^2 & -u^1 & 1 \\ -(u^2)^2 & u^2 & -1 \\ (u^3)^2 & -u^3 & 1 \end{bmatrix}.$$

Remark. If we multiply each row $\varphi_{(a)}^i$ of a Stäckel matrix by a function $f_i(q^i) \neq 0$, then we get a new Stäckel matrix $\tilde{\varphi}_{(a)}^i = f_i \varphi_{(a)}^i$ whose inverse $\tilde{\varphi}_{(a)}^i$ defines a new basis of the same KS-space. By multiplying the lines of (8.12) by ± 1 in a suitable way and by changing \mathbf{L} in $-\mathbf{L}$, we can get a Stäckel matrix which is of the Vandermonde type (see the case $n = 3$, for instance).

9. The functional independence of the eigenvalues of a L-tensor

The results of §8 hold without any assumption on the functional independence of the eigenvalues (u^i) of \mathbf{L} . Some of them may be constant. The only essential assumption is that they are pairwise and pointwise distinct. The following theorems deal with this matter.

Theorem 9.1. *If a torsionless CKT has functionally independent eigenvalues (u^i) , then it is a L-tensor (its eigenvalues are pointwise simple).*

Proof. In this case the eigenvalues define local orthogonal coordinates $q^i = u^i$ in which the tensor is diagonalized. We can apply theorem 6.2. Equations (6.2) become $\delta_i^k = (u^i - u^k)\partial_i \ln g^{kk} + 1$. For $i \neq k$ we have $u^i \neq u^k$. □

Note that in this case \mathbf{L} is a special type of L-tensor (the type considered by Crampin 2003 *a, b*).

Theorem 9.2. *Let \mathbf{L} be a L-tensor. (i) The eigenvalues (u^i) of \mathbf{L} are independent functions (i.e., they define locally an orthogonal coordinate system) if and only if \mathbf{L} is not invariant w.r.to a Killing vector \mathbf{X} . (ii) If there exists a Killing vector \mathbf{X} such that $[\mathbf{X}, \mathbf{L}] = 0$, then \mathbf{X} is a linear combination (with constant coefficients) of Killing vectors in involution which are eigenvectors of \mathbf{L} .*

In case (i) we have no *symmetry* of the separable web generated by the eigenvectors of \mathbf{L} . This means that all the structures associated with \mathbf{L} are not invariant w.r.to groups of isometries.

Proof. The eigenvalues of a (1,1) tensor \mathbf{L} are functionally independent, and thus they define a coordinate system, if and only if $\det[\partial_i u^j] \neq 0$. For a torsionless tensor with simple eigenvalues we have, w.r.to the associated coordinates, $\partial_i u^j = 0$ for $i \neq j$. Hence, the eigenvalues are independent functions if and only $\partial_i u^i \neq 0$ for all indices. (i) If an eigenvalue u^i of \mathbf{L} is constant, $\partial_i u^i = 0$, then from the expression (7.3) of the metric we see that the corresponding (normalized) coordinate q^i is ignorable. This means that $\partial/\partial q^i$ is a Killing vector which leaves invariant \mathbf{L} . Conversely, let us assume that there is a Killing vector \mathbf{X} such that $[\mathbf{X}, \mathbf{L}] = 0$. These two assumptions on \mathbf{X} are equivalent to $\{P_{\mathbf{X}}, P_{\mathbf{G}}\} = 0$ and $\{P_{\mathbf{X}}, P_{\mathbf{L}}\} = 0$. In an orthogonal coordinate system (q^i) associated with \mathbf{L} , these two equations read $\{X^i p_i, g^{jj} p_j^2\} = 0$ and $\{X^i p_i, u^j g^{jj} p_j^2\} = 0$, respectively. Since they are algebraic equations in p , to be satisfied for any value of these variables, they are equivalent to

$$\begin{cases} \sum_i X^i \partial_i g^{kk} - 2 g^{kk} \partial_k X^k = 0, & \left\{ \sum_i X^i \partial_i (u^k g^{kk}) - 2 u^k g^{kk} \partial_k X^k = 0, \right. \\ \left. g^{jj} \partial_j X^k + g^{kk} \partial_k X^j, \quad j \neq k, \right. & \left. \left\{ u^j g^{jj} \partial_j X^k + u^k g^{kk} \partial_k X^j, \quad j \neq k. \right. \right. \end{cases}$$

The first two equations characterize the Killing vectors in orthogonal coordinates. The second and the fourth equations imply $(u^j - u^k) g^{jj} \partial_j X^k = 0$ for $j \neq k$. Since $u^j \neq u^k$ we conclude that $\partial_j X^k = 0$ for $j \neq k$, which means that $X^i = X^i(q^i)$. Since $\partial_i u^k = 0$ for $i \neq k$, from the third equation it follows that $X^k \partial_k u^k g^{kk} + u^k \sum_i X^i \partial_i g^{kk} - 2 u^k g^{kk} \partial_k X^k = 0$. Due to the first equation (8.10), this last equation implies $X^k \partial_k u^k = 0$ (no summation over the index k). Up to a reordering of the coordinates, let us assume that $X^a = 0$ and $X^\alpha \neq 0$ for $a = 1, \dots, m$ and $\alpha = m+1, \dots, n$. From the last equation it follows that $u^\alpha = \text{const.}$ and $\mathbf{X} = X^\alpha \partial_\alpha$. At the beginning of this proof we have seen that if $u^\alpha = \text{const.}$, then q^α are ignorable coordinates (we always assume that the coordinates (q^i) are normalized so that the metric assume the form (7.3)). Thus, ∂_α are Killing vectors in involution and eigenvectors of \mathbf{L} . Since $\mathbf{X} = X^\alpha \partial_\alpha$ is a Killing vector, the components X^α must be constant. \square

Assume that the eigenvalues u^α ($u^\alpha = m+1, \dots, n$) of a L-tensor are constant and the remaining (u^a) ($a = 1, \dots, m$) are independent functions. Then we can choose associated orthogonal coordinates (q^i) = (q^a, q^α) such that $q^a = u^a$ and q^α are ignorable. From (7.1) it follows that

$$(u^a - u^b) \partial_a \ln g^{bb} + 1 = 0, \quad (u^a - u^\alpha) \partial_a \ln g^{\alpha\alpha} + 1 = 0,$$

being the remaining equations identically satisfied. Thus, we are faced with three cases: (I) $m = 0$, all $u^i = \text{const.}$ i.e, all q^i are ignorable: the manifold Q is locally flat, the coordinates (q^i) are orthogonal Cartesian coordinates, $g^{ii} = \text{const.}$ and \mathbf{L} is a constant tensor. (II) $0 < m < n$: in this case $g^{\alpha\alpha} \neq \text{const.}$ due to equation $(u^a - u^\alpha)\partial_a \ln g^{\alpha\alpha} + 1 = 0$. Condition $g^{\alpha\alpha} \neq \text{const.}$ means that the Killing vectors $\mathbf{X}_\alpha = \partial_\alpha$ are not translations (Eisenhart 1933, §52). (III) $m = n$, all eigenvalues are independent: this is the case examined in theorem 9.2

Taking into account the proof of the preceding theorem, cases (I) and (II) shows that:

Theorem 9.3. *Let \mathbf{L} be a L-tensor. (i) If \mathbf{L} has all constant eigenvalues, then the manifold Q is locally flat and $\mathbf{L} = \text{const.}$ (in the sense that all its components in Cartesian coordinates are constant). (ii) If \mathbf{L} is invariant w.r.to $m < n$ Killing vectors, then these vectors are not translations.*

10. L-potentials

In this section we propose a few remarks on the *potential functions* associated with a L-system. For further approaches to this matter we refer the reader to Ibort *et al.* 2000, Błaszak & Ma 2003. By (1.3) and (7.3) we have

Theorem 10.1. *A potential V is separable in a L-system if and only if, w.r.to normal coordinates (q^i) associated with \mathbf{L} ,*

$$\partial_i \partial_j V = \frac{1}{u^j - u^i} (\partial_j V - \partial_i V), \quad \partial_i = \frac{\partial}{\partial q^i}.$$

Note that in this theorem (as well as in the following) the eigenvalues u^i of \mathbf{L} may not be independent functions.

Theorem 10.2. *Let (\mathbf{K}_a) be the basis of the KS-space generated by a L-tensor \mathbf{L} according to formula (8.7). Then the functions $V_a(\underline{u}) \doteq \sigma_{a+1}(\underline{u})$ ($a = 0, 1, \dots, n-1$) fulfill equations $dV_a = \mathbf{K}_a dV$, with $V = V_0 = \text{tr}(\mathbf{L}) = \sum_i u^i$, and the functions $H_a = \frac{1}{2} P_{\mathbf{K}_a} + V_a$ are first integrals in involution.*

Proof. $\mathbf{K}_a dV = \sigma_a^i \partial_i V dq^i = \sigma_a^i dq^i = \partial_i \sigma_{a+1} dq^i = dV_a$. Apply (5.1). \square

11. L-pencils

There are Stäckel webs which are not L-webs. A necessary condition is the Robertson condition. However, also in manifolds where this condition is identically satisfied, for instance in constant curvature spaces, there are Stäckel systems which are not generated by a L-tensor. For instance, due to theorem 9.3, (ii), in a \mathbb{E}_n all translational webs (except the Cartesian web) are not L-webs. As remarked above, also the spherical-conical webs are not L-webs. This last case has suggested (Benenti 1992b) the introduction of a linear combination

$$\mathbf{L}(m) = \mathbf{L}_0 + m\mathbf{L}_1,$$

which is a L-tensor for all values the parameter $m \in \mathbb{R}$. We call **L-pencil** such an object.

Theorem 11.1. *Let $\mathbf{L} = \mathbf{L}_0 + m\mathbf{L}_1$ be a L-pencil. Then: (i) \mathbf{L}_0 has simple eigenvalues. (ii) \mathbf{L} is a CKT for all m if and only if \mathbf{L}_0 and \mathbf{L}_1 are CKT's. (iii) The condition $\mathbf{H}(\mathbf{L}) = 0$ is equivalent to*

$$\mathbf{H}(\mathbf{L}_0) = \mathbf{H}(\mathbf{L}_1) = 0, \quad L_{[i}^h \partial_{|h|} L_{j]}^k - L_{[i}^k \partial_{|i|} L_{j]}^h + L_{[i}^h \partial_{|h|} L_{j]}^k - L_{[i}^k \partial_{|i|} L_{j]}^h = 0.$$

(iv) *The Ricci tensor \mathbf{R} commutes with both \mathbf{L}_0 and \mathbf{L}_1 .*

The proof is straightforward. Let us apply the iterative formula (8.7) and theorem 8.7 to the tensor $\mathbf{L}(m)$. Since \mathbf{L} is polynomial of degree 1 in m , each tensor $\mathbf{K}_a(m)$ is at most of degree a : $\mathbf{K}_a(m) = \mathbf{H}_a m^a + \mathbf{H}_{a-1} m^{a-1} + \dots$

Theorem 11.2. (i) *The tensors $\mathbf{K}_a(0)$ form a KS-space. (ii) The tensors \mathbf{H}_a , given by the coefficients of maximal degree of $\mathbf{K}_a(m)$, if independent, form a KS-space. (iii) The Stäckel systems generated by these two KS-spaces satisfy the Robertson condition.*

Also for this statement we omit the proof. We remark that there are cases in which all the \mathbf{K}_a are of degree 1, as in the following example.

Example (Benenti 1992b): in \mathbb{E}_n we consider the tensor $\mathbf{L}(m) = \mathbf{L}_0 + m \mathbf{r} \otimes \mathbf{r}$. It represents (for $m \neq 0$) the planar moment of inertia of a massive body with total mass m and center of mass at the origin. \mathbf{L}_0 is a constant symmetric tensor (hence, a KT) with simple eigenvalues (a^i). $\mathbf{L}_1 = \mathbf{r} \otimes \mathbf{r}$ is a CKT whose eigenvalues are all zero except one ($= r^2$). It can be proved that: (i) \mathbf{L} is a L-pencil; (ii) the tensors $\mathbf{K}_a(0)$ form a KS-space corresponding to Cartesian coordinates; (iii) $\mathbf{K}_a(m)$ are all of degree 1 in m , $\mathbf{K}_a(m) = \mathbf{K}_a(0) + m \mathbf{H}_a$; (iv) the tensors \mathbf{H}_a form a KS-space corresponding to the conical spherical coordinates.

The above results stimulate investigations about the notion of L-pencil, also in relation with recent studies on the same concept and the notion of *cofactor pair system* (see e.g., Lundmark 2001, Rauch-Wojciechowski & Waksjö 2003, Marciniak & Błaszak 2002, Rauch-Wojciechowski, Marciniak & Lundmark 1999). A possible generalization is a **multi-pencil** of the type $\mathbf{L} = \mathbf{L}_0 + m^i \mathbf{L}_i$ with $(m^i) \in \mathbb{R}^k$ or $\mathbf{L} = m^i \mathbf{L}_i$ with $(m^i) \neq \mathbf{0}$. Basic examples are, in the Euclidean n -space, the case $\mathbf{L} = \mathbf{L}_0 + m \mathbf{r} \otimes \mathbf{r} + w \mathbf{c} \odot \mathbf{r}$ where \mathbf{c} is a constant unit vector, see (4.2). For lack of space, we cannot examine here which Stäckel webs in \mathbb{E}_n (or in $\mathbb{S}_n, \mathbb{H}_n$) are generated by a L-tensor or by a L-pencil.

12. The Riemannian background of the separation

Also for a better understanding of the orthogonal separation, it is necessary to study the Levi-Civita equations without any *a priori* assumptions on the separable coordinates q . The geometrical meaning of these equations has been firstly investigated by Kalnins & Miller 1980, 1981, and by Benenti 1980 (see also Benenti 1991, 1996). A result of these investigations in the following theorem (Benenti 1997):

Theorem 12.1. *The Hamilton-Jacobi equation $G + V = E$ is separable if and only if there exists a characteristic Killing pair (D, \mathbf{K}) such that*

$$DV = 0, \quad d(\mathbf{K} dV) = 0. \quad (12.1)$$

A **characteristic Killing pair** (D, \mathbf{K}) is made of a r -dimensional linear space D of commuting Killing vectors and a D -invariant Killing tensor \mathbf{K} having $m = n - r$ normal eigenvectors orthogonal to D and corresponding to distinct eigenvalues (N.B., these eigenvalues may not be simple). Since the elements of D are in involution, they generate r -dimensional orbits which are locally flat submanifolds. It can be shown that *if (D, \mathbf{K}) is a characteristic Killing pair, then D is normal*, in the sense that the distribution orthogonal to its elements is completely integrable. This means that there is a foliation of m -dimensional manifolds orthogonal to the orbits of D . The leaves of this foliation are isometric Riemannian manifolds (the isometries being generated by the orbits of D , assuming that they intersect the orbits in only one point).

With a characteristic Killing pair we associate **standard separable coordinates** $(q^i) = (q^a, q^\alpha)$ ($a = 1, \dots, m$, $\alpha = m + 1, \dots, n$) defined in this way: (i) dq^a are eigenforms of \mathbf{K} orthogonal to D (or, equivalently, the orbits of D are defined by equations $q^a = \text{const.}$); (ii) the (q^α) are the affine parameters, with zero value on an arbitrary m -dimensional orthogonal section \mathcal{Z} of the orbits of D , of Killing vectors \mathbf{X}_α forming a local basis of D . It follows that: (I) $\partial/\partial q^\alpha = \mathbf{X}_\alpha$ and the coordinates (q^α) are ignorable; (II) in these coordinates the contravariant metric tensor components assume the **standard form**

$$[g^{ij}] = \begin{bmatrix} [g^{aa}] & [0] \\ [0] & [g^{\alpha\beta}] \end{bmatrix}, \quad (12.2)$$

where $[g^{aa}]$ is a diagonal $m \times m$ matrix and $[g^{\alpha\beta}]$ is a $r \times r$ matrix, with $m = n - r$. Here we denote by $[0]$ zero-matrices of proper dimensions.

The rather long proof of this theorem is based on the following (Benenti 1980)

Theorem 12.2. *In an equivalence class of separable coordinates there exist coordinates in which the metric assume the standard form (12.2).*

Two separable systems are called **equivalent** if (in the intersection of their domains of definition) the corresponding separated solutions of the Hamilton-Jacobi equation generate the same Lagrangian foliation of T^*Q . The geometrical representation of an equivalent class is given by a **separable web**: it is a pair $(D, \mathcal{S}^a) = (D, (\mathcal{S}^1, \dots, \mathcal{S}^m))$, where D is a normal r -dimensional space of commuting KT's and (\mathcal{S}^a) is a family of m orthogonal foliations of submanifolds of codimension 1, all invariant w.r.to D (it follows that the orbits of D are the complete intersections of the (\mathcal{S}^a)). Moreover, the submanifolds (\mathcal{S}^a) are orthogonal to m eigenvectors of a D -invariant KT \mathbf{K} with distinct (but not necessarily simple) eigenvalues (so that (D, \mathbf{K}) is a characteristic Killing pair). It must be observed that the quotient set of the orbits of D is a m -dimensional manifold with a naturally induced orthogonal separable metric (g^{aa}) (it is isomorphic to any m -manifold orthogonal to D).

In this description of the separation we include the extreme cases: (I) $m = n$, $r = 0$, which corresponds to the an orthogonal separation.; (II) $m = 0$, $r = n$, which corresponds to the separation in Cartesian coordinates (in this case the manifold is flat). In Benenti *et al.* 2002a it is proved that

Theorem 12.3. *A characteristic Killing pair (D, \mathbf{K}) generates a m -dimensional space \mathcal{K}_m of Killing tensors with the following properties: (i) they are D -invariant and have m eigenvectors in common orthogonal to D ; (ii) they are in involution;*

(iii) if the **characteristic equations** (12.1) of a separable potential V are satisfied, then they are satisfied by all elements of \mathcal{K}_m .

Theorem 12.4. In a space \mathcal{K}_m having properties (i) and (ii), where D is a $n - m$ -space of Killing vectors in involution, all the common eigenvectors are normal.

Theorem 12.5. The functions $H_a = \frac{1}{2} P_{\mathbf{K}_a} + V_a$ and $P_{\mathbf{X}_\alpha} = X_\alpha^i p_i$, where $dV_a = \mathbf{K}_a dV$ and (\mathbf{X}_α) is a basis of D , form a complete system of integrals in involution.

Theorems 12.2 and 12.3 show that if D admits an orthogonal basis, $\mathbf{X}_\alpha \cdot \mathbf{X}_\beta = 0$ for $\alpha \neq \beta$, then the standard coordinates are orthogonal. This means that the separation occurs in orthogonal coordinates. It can be proved by a coordinate-independent method that (Benenti 1992a)

Theorem 12.6. On a manifold with constant curvature any normal space D of Killing vectors in involution has an orthogonal basis.

As a consequence,

Theorem 12.7. On a manifold of constant curvature the geodesic separation always occurs in orthogonal coordinates.

This important property was discovered by Kalnins & Miller (1984, 1986) (see also Kalnins 1986), within a coordinate-dependent approach. This theorem can be extended to a natural Hamiltonian $H = G + V$.

References

- Ankiewicz, A. & Pask, C. 1983 The complete Whittaker theorem for two-dimensional integrable systems and its applications. *J. Phys. A: Math. Gen.* **16**, 4203.
- Benenti, S. 1980 Separability structures on Riemannian Manifolds. In *Lecture Notes in Mathematic* **863**, 512-538.
- Benenti, S. 1991 Separation of variables in the geodesic Hamilton-Jacobi equation. In *Progress in Mathematics* **99**, 1-36.
- Benenti, S. 1992a Orthogonal separation of variables on manifolds with constant curvature. *Diff. Geom. Appl.* **2**, 351-367.
- Benenti, S. 1992b Inertia tensors and Stäckel systems in the Euclidean spaces. *Rend. Semin. Mat. Univ. Polit. Torino* **50**, 315-341.
- Benenti, S. 1993 Orthogonal separable dynamical systems. In *Proceedings of the 5th International Conference on Differential Geometry and Its Applications, Silesian University at Opava, August 24-28, 1992, O.Kowalski and D.Krupka Eds. Differential Geometry and Its Applications* **1**, 163-184. Web edition: ELibEMS, <http://www.emis.de/proceedings>.
- Benenti, S. 1996 Connections and Hamiltonian Mechanics. In *Gravitation Electromagnetism and Geometrical Structures*, Ferrarese, G. Ed., pp. 185-206. Bologna: Pitagora Editrice.
- Benenti, S. 1997 Intrinsic characterization of the variable separation in the Hamilton-Jacobi equation. *J. Math. Phys.* **38** (12), 6578-6602.
- Benenti, S., Chanu, C. & Rastelli, G. 2000 The super-separability of the three-body inverse-square Calogero system. *J. Math. Phys.* **41** (7), 4654-4678.
- Benenti, S., Chanu, C. & Rastelli, G. 2001 Variable separation for natural Hamiltonians with scalar and vector potentials on Riemannian manifolds. *J. Math. Phys.* **42** (5), 2065-2091.

- Benenti, S., Chanu, C. & Rastelli, G. 2002a Remarks on the connection between the additive separation of the Hamilton-Jacobi equation and the multiplicative separation of the Schrödinger equation. I. The completeness and Robertson conditions. *J. Math. Phys.* **43** (11), 5183-5222.
- Benenti, S., Chanu, C. & Rastelli, G. 2002b Remarks on the connection between the additive separation of the Hamilton-Jacobi equation and the multiplicative separation of the Schrödinger equation. II. First integrals and symmetry operators. *J. Math. Phys.* **43** (11), 5223-5253.
- Błaszak, M. & Ma, W.-X. 2003 Separable Hamiltonian equations on Riemann manifolds and related integrable hydrodynamic systems. *J. Geom. Phys.* **47**, 21-42.
- Błaszak, M. 2003 Separability theory of Gel'fand-Zakharevic systems on Riemannian manifolds. Preprint, A. Mickiewics University, Poznań.
- Błaszak, M. & Badowski, L. 2003 From separable geodesic motion to bihamiltonian dispersionless chains. Preprint, A. Mickiewics University, Poznań.
- Bolsinov, A. V. & Matveev V. S. 2003 Geometrical interpretation of Benenti systems. *J. Geom. Phys.* **44**, 489-506.
- Boyer, C. P., Kalnins, E. G. & Miller Jr., W. 1986 Stäckel-equivalent integrable Hamiltonian systems. *SIAM J. Math. Anal.* **17**, 778-797.
- Bruce, A.T., McLenaghan, R. & Smirnov, R.G. 2001 Benenti's theorem and the method of moving frames. *Rep. Math. Phys.* **48**, 227-234.
- Crampin, M. 2003a Conformal Killing tensors with vanishing torsion and the separation of variables in the Hamilton-Jacobi equation. *Differential Geom. Appl.* **18**, 87-102.
- Crampin, M. 2003b Projectively Equivalent Riemannian Spaces as Quasi-bi-Hamiltonian Systems. *Acta Applicandae Mathematicae* **77** (3), 237-248.
- Deeley, J., Horwood, J. T., McLenaghan, R. & Smirnov, G.R. 2004 Theory of algebraic invariants of vector spaces of Killing tensors: Methods of computing the fundamental invariants. *Proceedings of Inst. Of Math. of NAS of Ukraine*, to appear.
- Eisenhart, L. P. 1933 *Continuous groups of transformations*. Princeton: Princeton University Press.
- Eisenhart, L. P. 1934 Separable systems of Stäckel. *Ann. Math.* **35**, 284-305.
- Eisenhart, L. P. 1949 *Riemannian geometry*. Princeton: Princeton University Press.
- Fettis, H. E. 1950 A method for obtaining the characteristic equation of a matrix and computing the associated modal columns. *Quart. Appl. Math.* **8**, 206-212.
- Haantjes, J. 1955 On X_m -forming sets of eigenvectors. *Proc. Kon. Ned. Ak. Wet. Amsterdam A* **58** (2), 158-162.
- Kalnins, E. G. 1986 *Separation of Variables for Riemannian Spaces of Constant Curvature*. Pitman Monographs **28**. Harlow: Longman Scientific & Technical.
- Kalnins, E. G. & Miller Jr., W. 1980 Killing tensors and variable separation for Hamilton-Jacobi and Helmholtz equations. *SIAM J. Math. Anal.* **11**, 1011-1026.
- Kalnins, E. G. & Miller Jr., W. 1981 Killing tensors and nonorthogonal variable separation for Hamilton-Jacobi equations. *SIAM J. Math. Anal.* **12**, 617-639.
- Kalnins, E. G. & Miller Jr., W. 1983 Intrinsic characterization of variable separation for the partial differential equations of Mechanics. In *Proceedings of IUTAM-ISIMM Symposium on Modern Developments in Analytical Mechanics, Torino 1982, Atti Accad. Sci. Torino* **117**, Vol.2, 511-533.
- Kalnins, E. G. & Miller Jr., W. 1984a Separation of variables on n -dimensional Riemannian manifolds - II. The n -dimensional hyperboloid \mathbf{H}_n . University of Waikato Research Report **103**.
- Kalnins, E. G. & Miller Jr., W. 1984b Separation of variables on n -dimensional Riemannian manifolds -III. Conformally Euclidean spaces \mathbf{C}_n . University of Waikato Research Report **105**.

- Kalnins, E. G. & Miller Jr., W. 1986 Separation of variables on n -dimensional Riemannian manifolds - I. The n -sphere \mathbf{S}_n and Euclidean n -space \mathbf{R}_n . *J. Math. Phys.* **27**, 1721-1736.
- Ibort, A., Magri, F. & Marmo, G. 2000 Bihamiltonian structures and Stäckel separability. *J. Geom. Phys.* **33**, 210-228.
- Levi-Civita, T. 1896 Sulle trasformazioni delle equazioni dinamiche. *Ann. di Matem.* **24**, 255-300.
- Levi-Civita, T. 1904 Sulla integrazione della equazione di Hamilton-Jacobi per separazione di variabili. *Math. Ann.* **59**, 383-397.
- Lundmark, H. 2001 A new class of integrable Newton systems. *J. Nonlin. Math. Phys.* **8**, supplement, 195-199.
- Marciniak, K. & Błaszak, M. 2002 Separation of variables in quasi-potential systems of bi-cofactor form. *J. Phys. A: Math. Gen.* **35**, 2947-2964.
- Marshall, I. & Wojciechowski, S. 1988 When is a Hamiltonian system integrable?. *J. Math. Phys.* **29**, 1338-1346.
- McLenaghan, R. & Smirnov, R.G. 2002 Intrinsic Characterizations of Orthogonal Separability for Natural Hamiltonians with Scalar Potentials on Pseudo-Riemannian Spaces. *J. Nonlin. Math. Phys.* **9**, Suppl. 1, 140-151.
- Miller Jr., W. 1988 Mechanism for variable separation in partial differential equations and their relationship to group theory. In *Symmetry and nonlinear phenomena*. Singapore: World Scientific.
- Moser, J. 1981 *Integrable Hamiltonian Systems and Spectral Theory*. Lezioni fermiane. Accademia Nazionale dei Lincei, Scuola Normale Superiore di Pisa.
- Nijenhuis, A. 1951 X_{n-1} -forming sets of eigenvectors. *Nederl. Akad. Wetensch. Proc.* **54A**, 200-212.
- Rauch-Wojciechowski, S. 2004 From Jacobi problem of separation of variables to theory of quasipotential Newton equations. *This volume*.
- Rauch-Wojciechowski, S. & Waksjö, C. 2003 Stäckel separability for Newton systems of cofactor type. Preprint, University of Linköping, Linköping.
- Rauch-Wojciechowski S., Marciniak, K. & Lundmark, H. 1999 Quasi-Lagrangian systems of Newton equations. *J. Math. Phys.* **40**(12), 6366-6398.
- Schouten, J. A. 1954 *Ricci Calculus*. Berlin: Springer.
- Shapovalov, V. N. 1981 Stäckel spaces. *Siberian Math. J.* **20**, 790-800.
- Souriau, J.M. 1950 Le calcul spinoriel et ses applications. *Recherche Aronautique* **14**, 3-8.
- Stäckel, P. 1893 Über die Bewegung eines Punktes in einer n -fachen Mannigfaltigkeit. *Mat. Ann.* **42**, 537-563.
- Stäckel, P. 1897 Über quadatische Integrale der Differentialgleichungen der Dynamik. *Ann. Mat. Pura Appl.* **26**, 55-60.
- Waksjö, C. 2000 Stäckel Multipliers in Euclidean Spaces. *Linköping Studies in Sciences and Technology. Theses* **833**.
- Waksjö, C. & Rauch-Wojciechowski, S. 2003 How to find separation coordinates for the Hamilton-Jacobi equation: a criterion of separability for natural Hamiltonian systems. *Math. Phys. Anal. Geom.* To appear.
- Whittaker, E.T. 1937 *A treatise on the analytical dynamics of particles and rigid bodies*. Cambridge: Cambridge Univ. Press.
- Woodhouse, N. M. J. 1975 Killing tensors and the separation of the Hamilton-Jacobi equation. *Commun. Math. Phys.* **44**, 9-38.